

Medication Guide Prior to Endoscopy Procedure

DO NOT HOLD – Continue These Medications	HOLD – Medications to Stop	When to Stop (Hold Time)
<p><u>Beta-blockers</u> -Metoprolol, Labetolol, Carvedilol, Propanolol, Atenolol</p> <p><u>Calcium-Channel Blockers</u> -Amlodipine, Diltiazem, Nefedipine, Verapamil</p> <p><u>Anti-hypertensives/ Heart Failure</u> -Lisinopril, Enalapril, Captopril, Benzapril -Hydralazine, Isosorbide, Irbesartan -Entresto, Minoxidil</p> <p><u>Anti-Arrhythmics</u> -Amiodarone, Digoxin, Flecainide, Disopyramide, Procainamide</p> <p><u>Diuretics</u> Furosemide, Hydrochlorothiazide, Bumetanide, Diovan, Lotensin, Torsemide, Spironolactone</p> <p><u>Opioids</u> -Fentanyl patch, Codeine, Tramadol, MS Contin, Hydromorphone, Methadone)</p> <p><u>Acid Reflux</u> -Famotidine, Pantoprazole, Ranitidine, Omeprazole -Oxetidine</p> <p><u>Steroids</u> -Prednisone, Hydrocortisone</p> <p><u>Anti-depressants/ Parkinson's</u> -Risperidone, Haloperidol (Haldol), Clozapine, Clonidine, Methyldopa, Bupropion -Lithium, Valproic Acid, Lamotrigine, Topiramate -Carbidopa-levodopa, Amantidine, Pramipexole, Entacapone -Trazadone</p> <p><u>Other Medications</u> -Levetiracetam (Keppra), Gabapentin, Phenytoin -Lorazepam, Diazepam -Levothyroxine -Albuterol, Ipratropium, Fluticasone -Statins (Cholesterol) -Estradiol, Tamoxifen -Doxazosin, Tamsulosin, Prazosin, Finasteride</p>	<p><u>GLP-1</u> - Ozempic (Semaglutide), Wegovy, Trulicity (Dulaglutide), Mounjaro (Tirzepatide), Byetta (Exenatide)</p>	Hold One week before procedure
	<p>- Repaglinide, Tanzeum, Rybelsus, Exenatide, Liraglutide, Lixisenatide</p>	Hold 24 hours before procedure
	<p><u>SGL-2</u> - Farxiga, Jardiance, Synjardy, Brenzavvy, Dapagliflozin, Empagliflozin, Bexagliflozin, Glyxambi, Trijardy, Invokana</p>	Hold for 3 days before procedure
	<p>-Steglatro (Ertugliflozin), Steglujan</p>	Hold for 4 days before procedure
	<p><u>Other Diabetic Medications</u> -Januvia (Sitagliptin), Tradjenta (Linagliptin), Onglyza (Saxagliptin)</p>	Hold for 2 days before procedure
	<p>-Nesina (Alogliptin), Galvus (Vildagliptin)</p>	Do not take the day of procedure
	<p>-Glipizide, Glyburide, Glimepiride, Repaglinide, Mateglinide</p>	Do not take the day of procedure
	<p>-Acarbose, Glyset (Miglitol), Metformin</p>	Do not take the day of procedure
	<p><u>Anticoagulants / Antiplatelets</u> -Eliquis (Apixaban), Xarelto (Rivaroxaban), Pradaxa (Dabigatran), Savaysa/ Lixiana (Edoxaban)</p>	Hold for 2 days before procedure
	<p>-Coumadin/ Jantoven (Warfarin),</p>	Hold 5-7 days before procedure
	<p>-Plavix (Clopidogrel), Effient (Prasugrel), Pletal (Cilostazol), Brilinta (Ticagrelor)</p>	Hold 5-7 days before procedure
	<p>-Aggrenox (Aspirin/dipyridamole)</p>	Hold for 2 days before procedure
	<p>-Trental (Pentoxifylline)</p>	Hold for 2 days before procedure
	<p>-Lovenox (Enoxaparin) injections</p>	Hold 24 hours before procedure
	<p><u>Other Medications</u> -Adipex-P (Phentermine), Lomaira (Phentermine 8 mg), Qsymia (Phentermine/ Topiramate)</p>	Hold 2 weeks before procedure
	<p>-Contrave (Bupropion/ Naltrexone)</p>	Hold 3 days before procedure
	<p>-Vyvanse (Lisdexamfetamine), Amphetamine, Adderall (Amphetamine/ Dextroamphetamine), Bupropion **If being used for weight loss**</p>	Hold 3 days before procedure
	<p>-Ergotamine (Ergomar, Cafergot)</p>	Hold 2 days before procedure